



Legislation Details (With Text)

File #: Res 0691-2011 **Version:** * **Name:** NYC Dept of Health and Mental Hygiene to study the effects of fluoride in tap water.

Type: Resolution **Status:** Filed

In control: Committee on Health

On agenda: 3/2/2011

Enactment date: **Enactment #:**

Title: Resolution calling upon the New York City Department of Health and Mental Hygiene to study the effects of fluoride in tap water.

Sponsors:

Indexes:

Attachments:

Date	Ver.	Action By	Action	Result
3/2/2011	*	City Council	Introduced by Council	
3/2/2011	*	City Council	Referred to Comm by Council	
12/31/2013	*	City Council	Filed (End of Session)	

Res. No. 691

Resolution calling upon the New York City Department of Health and Mental Hygiene to study the effects of fluoride in tap water.

By Council Members James, Cabrera, Chin, Mealy, Mendez, Rose, Williams and Koo

Whereas, Water fluoridation is the practice of adding fluoride compounds to water with the intended purpose of reducing tooth decay; and

Whereas, Fluoridation of drinking water has been common in the United States for more than fifty years, and most of the country's municipalities fluoridate their water supplies; and

Whereas, In 1986, the Environmental Protection Agency (EPA), established a maximum allowable concentration of 4 milligrams of fluoride per liter of water (4mg/L), a guideline designed to prevent the public from being exposed to harmful levels of fluoride; and

Whereas, According to a 2002 study, about two-thirds of the U.S. population and 46 of the nation's 50 largest cities receive fluoride through their community water system; and

Whereas, The Centers for Disease Control and Prevention (CDC) have proclaimed community water fluoridation (along with vaccinations and infectious disease control) as one of ten great public health achievements of the 20th century; and

Whereas, According to the American Dental Association (ADA), water fluoridation reduces dental decay by 40 to 59 percent, and past comprehensive reviews of the safety and effectiveness of fluoride in water have concluded that water fluoridation is safe and is the most cost-effective way to prevent tooth decay among populations living in areas with adequate community water supply systems; and

Whereas, Concerns have been raised, however, over the quality of previous research demonstrating the efficacy and safety of water fluoridation, as growing evidence suggests that fluorodiation poses serious health risks affecting teeth, bones, the brain and the thyroid gland; and

Whereas, In 2005, the U.S. National Institutes of Health (NIH), a research agency associated with the U.S. Department of Health and Human Services, evaluated 562 dental studies on fluoride use and reported that the studies were small, poorly described, or otherwise methodologically flawed; and

Whereas, In March 2006, the National Research Council (NRC) conducted a study to determine whether the current amount of fluoride allowed in drinking water poses a health risk to Americans; and

Whereas, After reviewing research on various health effects from exposure to fluoride, including studies conducted in the last 10 years, the NRC reported that constant exposure to fluoride at the current maximum level results in severe dental fluorosis (tooth enamel loss and pitting) in children, and increases the risk of bone fractures and skeletal fluorosis (painful stiffening of the joints) in adults; and

Whereas, The NRC concluded that the EPA's current limit for fluoride in drinking water does not protect against adverse health effects, and recommended that the federal government lower its fluoridation level limit because of health risks to both children and adults, but did not indicate what the limit should be; and

Whereas, Optimal fluoride levels for drinking water, as recommended by the U.S. Public Health Service and the CDC, range from 0.7 mg/L for warmer climates to 1.2 mg/L for cooler climates to account for the

tendency for people to drink more water in warmer climates; and

Whereas, All New York City tap water has been fluoridated since 1966 in accordance with the New York City Health Code, at a concentration of one milligram per liter (1mg/L); and

Whereas, Some researchers question whether even adding 1mg/L of fluoride into tap water is acceptable since there is no universally accepted optimal level for daily fluoride intake and there lacks scientifically valid evidence proving the safety or effectiveness of water fluoridation; and

Whereas, An article published in the *Journal of Public Health Dentistry* in 2003 stated that despite fluoridation, severe tooth decay is responsible for two-thirds of hospital visits by children under six in New York State; and

Whereas, According to a report by the CDC, proportionately more children in New York City required cavity-related hospitalizations than two of New York State's largest non-fluoridated counties, Suffolk and Nassau (in Long Island); and

Whereas, Similarly, twenty-one percent of Brooklyn's and twenty percent of Queens' residents have less teeth as compared to residents of non-fluoridated Suffolk and Nassau counties; and

Whereas, Although most fluoridated water contains much less than the EPA limit, studies challenging the safety and effectiveness of fluoridation have raised uncertainty about whether these lower amounts help to prevent health risks; and

Whereas, As there seems to be a significant amount of questionable data on the safety and efficacy of fluoride in tap water, additional extensive studies should be conducted to determine any actual effects; now, therefore, be it

Resolved, That the Council of the City of New York calls upon the Department of Health and Mental Hygiene to study the effects of fluoride in tap water.

Res No. 1392-2008
LS #2096
2/23/11